

Embryonic development of the proepicardium and coronary vessels

ANNA RATAJSKA^{1,*}, ELZBIETA CZARNOWSKA² and BOGDAN CISZEK³

¹Department of Pathological Anatomy, Medical University of Warsaw, ²Department of Pathology, Childrens' Memorial Health Institute and ³Department of Anatomy, Medical University of Warsaw, Poland

ABSTRACT In the last few years, an increasing interest in progenitor cells has been noted. These cells are a source of undifferentiated elements from which cellular components of tissues and organs develop. Such progenitor tissue delivering stem cells for cardiac development is the proepicardium. The proepicardium is a transient organ which occurs near the venous pole of the embryonic heart and protrudes to the pericardial cavity. The proepicardium is a source of the epicardial epithelium delivering cellular components of vascular wall and interstitial tissue fibroblasts. It contributes partially to a fibrous tissue skeleton of the heart. Epicardial derived cells play also an inductive role in differentiation of cardiac myocytes into conductive tissue of the heart. Coronary vessel formation proceeds by vasculogenesis and angiogenesis. The first tubules are formed from blood islands which subsequently coalesce forming the primitive vascular plexus. Coronary arteries are formed by directional growth of vascular protrusions towards the aorta and establishing contact with the aortic wall. The coronary vascular wall matures by attaching smooth muscle cell precursors and fibroblast precursors to the endothelial cell wall. The cells of tunica media differentiate subsequently into vascular smooth muscle by acquiring specific contractile and cytoskeletal markers of smooth muscle cells in a proximal - distal direction. The coronary artery wall matures first before cardiac veins. Maturity of the vessel wall is demonstrated by the specific shape of the internal surface of the vascular wall.

KEY WORDS: *epicardium, hematopoiesis, vasculogenesis, coronary vessel development, mesenchymal cell*

Introduction

The tissues and organs of multicellular organisms develop from the pool of undifferentiated progenitor cells. The example of such a pool of progenitor cells is vertebrate proepicardium, which contains cells for the development of the heart. The proepicardium is a transient organ which is located near the venous pole of the embryonic heart and protrudes to the pericardial cavity (Virágh *et al.*, 1993). The current data on coronary vessel development indicate that the proepicardium is a source of the epicardial epithelium (Männer *et al.*, 2001), which participates in the formation of the vascular wall containing endothelial cells, smooth muscle cells and fibroblasts (Männer *et al.*, 2001). The proepicardium is also a source of cells forming connective tissue of the heart valves and modulating formation of the heart skeleton (Winter, Gittenberger-de Groot 2007). The proepicardium is described as the "organ" because it is a source of a heterogenous

population of progenitor cells, which participate in the development of various structures of the heart. These cells participate in such processes as vasculogenesis, angiogenesis, morphogenesis and remodeling of extracellular matrix.

Formation of the proepicardium

The primordium of the proepicardium is found on the surface of the septum transversum in mammals, and on the surface of the sinus venosus near the embryonic liver in birds. The proepicardium primordium forms as the multicellular protrusion of the pericardial serosa covering both horns of the sinus venosus (in mammals) or the right sinus horn (in birds) (Männer *et al.*, 2001, Schulte *et al.*, 2007). These protrusions contain hyaluronic acid and a small amount of fibronectin (Kálmán *et al.*, 1995; Männer *et al.*, 2001). At this stage of development, in mammals, the embryonic liver grows into the septum transversum. In birds, which lack the

*Address correspondence to: Anna Ratajska, Department of Pathological Anatomy, The Medical University of Warsaw, Chalubinskiego 5, 02-004 Warsaw, Poland. Fax: +48-(22)-629-9892. e-mail: arataj@ib.amwaw.edu.pl




Fig. 1. The proepicardium of a mouse embryo at stage 9.5 dpc. (A) A sagittal section of the embryo showing the location and the structure of the proepicardium (arrow), mag. 50x. (B) The proepicardium releasing a vesicle (arrow), mag. 90x. (C) Transmission electron microscopy image of the proepicardial surface with epithelial-like cells (arrow), mag. 2600x. (D) Transmission electron microscopy image of the proepicardium with loosely arranged mesenchymal cells; mag. 3300x; a, atrium; v, ventricle.

diaphragm and in which the septum transversum does not develop, the proepicardium forms near the embryonic liver. Thus, the protrusion forms in the pericardial cavity, near the septum transversum (in mammals) or fetal liver (in birds). It resembles cauliflower (in birds) and grapes (in mammals) (Ho and Shimada 1978; Virágh, Challice 1981). These protrusions give rise to the proepicardium. In the next step of proepicardium development these protrusions grow within the pericardial cavity (Fig. 1A). In birds such as quail or chicken the proepicardium reaches about 200µm in diameter, in mice - about 60-80 µm and is correspondingly larger in human. It reaches its maximum diameter at the 4-5 week of the prenatal life in human, at 9.5 dpc in mice and at HH15-17 in birds (Hamburger and Hamilton 1951).

Polish scientist Kurkiewicz was the first who described the proepicardial protrusions (he called them villi), and suggested that they give rise to the epicardium (Kurkiewicz 1909). The proepicardium is covered with embryonic epithelial cells while the core of proepicardium consists of undifferentiated mesenchymal cells, loosely arranged and connected by intercellular junctions (Fig. 1 B,C). Further development of the proepicardium after it

reaches its maximum size differs in mammals and in birds but in both cases, proepicardium differentiates into the epicardium (Virágh, Challice 1981; Hiruma, Hirakow 1989).

Immunohistochemical and genetic markers of the proepicardium

Early proepicardium does not express any markers of differentiated endothelial cells or their precursors such as angioblast (CD34, PECAM-1, VEGFR1, VEGFR2), fibroblast, hematopoietic cells (CD45, CD34), smooth muscle cells (alpha-actin) or other markers (Fig. 2). Endothelial cells are detected in the proepicardium in later stages of development when the proepicardium reaches the heart surface and attaches to it (Kattan *et al.*, 2004; Guadix *et al.*, 2006).

On the other hand the proepicardium contains the transcription factor of Wilm's tumor (WT-1) suppressor gene, retinaldehyde dehydrogenase2 (RALDH2) (Pérez-Pomares *et al.*, 2002b), and *Tbx* (Hatcher *et al.*, 2004) and *Cfcb* gene products (Schlueter *et al.*, 2006). Mutant mice for *WT-1* display abnormalities in the epicardium formation with a reduced number of subepicardial mesenchymal cells (Moore *et al.*, 1999). Normal development of the proepicardium requires expression of BMP factor (Schlueter *et al.*, 2006) and GATA4 transcription factor (Watt *et al.*, 2004).

Formation of the epicardium

In embryonic stages of HH14 and HH17 in birds (and at 9-10 dpc in mice) through further bending of the head region of the embryo and concomitant heart looping the dorsal surface of the heart loop approaches the ventral wall of the sinus venosus. In birds, the growing protrusions of the proepicardium approach the dorsal surface of the heart and attach to it. This forms a bridge in the pericardial cavity, between the ventral surface of the sinus venosus and dorsal wall of the heart (Männer 1993). In birds this bridge is composed of extracellular matrix (ECM) including heparan sulfate and fibronectin (Ho, Shimada 1978; Männer 1992; Männer 1993; Virágh *et al.*, 1993; Männer 1999; Nahirney *et al.*, 2003). The bridge was also observed in mammals (Nesbitt *et al.*, 2006) and its composition is similar. Looping heart is covered by cells deriving from the proepicardium, which form a simple squamous epithelium i.e. the epicardium. Initially these cells adhere to the dorsal surface of the heart. Then cells of the epicardium proliferate and spread on the heart surface (Hiruma, Hirakow 1989).

In mammals, formation of the epicardium from the proepicardium, occurs via a release of free-floating vesicles into the pericardial cavity. These vesicles are released from the tip of the proepicardium (Virágh, Challice 1981; Kuhn, Liebherr 1988; Van den Eijnde *et al.*, 1995). In mice, the highest number of free-floating vesicles in the pericardial cavity occurs at 9.5-10 dpc. These vesicles reach also the dorsal surface of the heart, attach to it, and subsequently flatten and spread on the surface of the "naked" heart. Expansion of the epicardium occurs by proliferation of proepicardial and mesothelial cells. (Fig. 3). These cells express cytokeratins, which are the marker of epithelial cells (Vrancken Peeters *et al.*, 1995). There is a distinct spatial-temporal pattern of the spreading of the mesothelial cells on the surface of the heart. The apical surfaces of the proepicardial clusters bleb off from the proepicardium, attach to the dorsal wall

of the atrioventricular sulcus and become the epicardial mesothelium. Subsequently the epicardium covers almost the whole dorsal surface of the ventricle moving towards the atrium and bulbus cordis, with a delay in the covering of the ventral surface. Beginning from the dorsal surface of the heart, the epithelial cells form the rings in the atrioventricular and ventriculo-bulbar sulcuses, subsequently cells spread to the right and to the left around the looping heart on the ventricular surface (the heart is U-shaped at this stage of development). Then, cells spread cranially and caudally reaching the outflow tract and atria (the shape of the heart resembles an "S") (Hiruma, Hirakow 1989; Vrancken Peeters *et al.*, 1995). The front line of the spreading epicardial cells covering the looping heart has a semilunar shape. The cells expand on the heart surface radially from the atrioventricular sulcus. In mice, by the end of 11 dpc, the heart is entirely covered by the epicardium. This corresponds to the stage HH26 in birds. In chicken, covering the heart with the epicardium takes 2 days.

All proepicardial cells transform into the cells of epicardium. The proepicardium disappears by the end of the 5th week of human development (in mice by 10.5-11 dpc). The limited release of cells is observed in a group of mesothelial cells from the dorsal surface of the pericardial cavity in several animal species (Männer *et al.*, 2001). We noticed such cell release in the pericardium of mouse embryos (Fig. 4).

The epithelial-mesenchymal transformation of epicardial cells

In further stages of development the epicardium forms cellular components of the subepicardium. Subepicardium synthesizes extra cellular matrix built from glycosaminoglycans, collagen, elastin, and fibronectin. This occurs through the invagination of the subpopulation of epicardial cells under the surface of the epicardium. The invaginating cells lose their epithelial character and transform into undifferentiated mesenchymal cells (Dettman *et al.*, 1998; Pérez-Pomares *et al.*, 1998). This phenomenon is called the epithelial-mesenchymal transformation (EMT). It is accompanied by expression of the Slug and Snail transcription factors, whose presence are necessary for this process (Carmona *et al.*, 2000; Cano *et al.*, 2000). Other growth factors like TGF- β 1 and -2 also stimulate this process through their constitutive





Fig. 4. Mouse embryo at the stage of 9.5 dpc. The dorsal surface of the peritoneal-pericardial cavity with released epithelial cells, which form vesicle-like structures similar to those observed in the proepicardium. (A) The area in the rectangle is enlarged in (B,C,D). (B,C,D) Serial sections; v, ventricle; magnifications: (A) 60x, (B-D) 220x.

receptor- ALK2 (Olivey *et al.*, 2006). Mesenchymal cells are a source of extracellular matrix. Erythrocytes and angioblasts are found in newly formed subepicardium (Virágh *et al.*, 1993; Ratajska *et al.*, 2006, Tomanek *et al.*, 2006a). Erythrocytes are surrounded by angioblasts forming the primitive vascular vesicles (Fig. 5).

Similar epithelial-mesenchymal transformation occurs within the endocardial cushion tissue of the heart; here, the endocardial endothelial cells lose their epithelial character and transform into the mesenchymal cells of the cushion tissue (Markwald *et al.*, 1977).

The mesenchymal cells which derive from the epicardium (epicardial-derived mesenchymal cells - EPDC) build a layer of connective tissue, which is the thickest in the atrioventricular and interventricular sulcuses. This connective tissue will form the epicardium, which will be covered with epithelial cells and filled with fibroblasts and extracellular matrix built from collagen, glycosaminoglycans, elastic and collagen fibers. Subepicardial space is occupied by primitive vascular vesicles, which are not yet connected to the systemic circulation (Kálmán *et al.*, 1995).

The routs of migration of mesenchymal cells derived from the epicardium (and earlier from the proepicardium) are known from the analysis of interspecies chimeras, in which the quail

Fig. 2 (Left). The sagittal section of the proepicardium of mouse embryo at the stage of 9.5 dpc. (A) A histological section of the proepicardium (arrow) stained with hematoxylin-eosin; numerous vesicles released to the pericardial cavity are visible. (B) Section stained with anti-CD34 antibodies; CD34 positive cells are located in the dorsal aorta and at the base of the proepicardium; v, ventricle; da, dorsal aorta. Magnification: 200x.

Fig. 3 (Right). The heart of mouse stage of 10 dpc embryo partially covered with the epicardium. (A) The area covered with the epicardium is located between arrows; mag. 200x. (B) Proliferating epicardial cell (arrow), mag. 380x.

proepicardium was inserted into the chicken pericardial cavity (Gittenberger-de Groot *et al.*, 1998). The presence of quail specific nuclear marker allowed tracking the migration of quail cells in the chimeric chicken embryo. In addition, the retroviral tagging of the proepicardium with a reporter gene for β -galactosidase allowed tracking the migration of proepicardium-derived cells (Mikawa, Fishman 1992; Mikawa, Gourdie 1996).

It is not known whether there are separate populations of mesothelial cells able to differentiate into different cell types: endothelial, smooth muscle or fibroblast, or whether all EPDC are able to differentiate into all types of cells, depending on the action of specific growth factors (Dettman *et al.*, 1998; Pérez-Pomares *et al.*, 2002a). *In vitro* studies showed that the growth factors are able to induce all these transformations in certain populations of mesenchymal cells. Also the epicardial epithelium has a great capacity to differentiate into other cell types (Wessels, Pérez-Pomares 2004). The process begins with initial dedifferentiation of flat epithelial cells into mesenchymal cells. Although the proepicardium does not have differentiated cells it has an enormous potential to differentiate into vascular cells *in vivo* and *in vitro* (Guadix *et al.*, 2006, Pérez-Pomares *et al.*, 2006).

Knockout mice devoid of VCAM-1 adhesive molecule do not develop the epicardium and their myocardium is very thin. A large amount of blood cells is present within the pericardial cavity of these knockout mice (Kwee *et al.*, 1995). Photoablation experiments on chicken hearts showed that the hearts are devoid of the proepicardium and epicardium, are thin-walled and aneurysmal, have defects of coronary vasculature and develop hemo- and hydropericardium (Männer *et al.*, 2005). This indicates that the proepicardium plays a role in maintaining the mechanical properties of the heart wall.

Formation of coronary vessels

The coronary vessels develop from the blood islands which are the aggregates of endothelial cells and erythrocytes and are not connected to the systemic circulation (Hirakow 1983; Rongish *et al.*, 1994; Ratajska, Fiejka 1999). The blood islands coalesce to form the primitive vessels. In birds, at the HH23 stage, the blood islands, erythrocytes and capillary vessels are most numerous within the subepicardial space (Hiruma, Hirakow 1989) and the vascularization of the heart is preceded by the development of the epicardial cover. In birds and mammals, the earliest vascular structures are visible within the sinus venosus and the dorsal part of the atrioventricular sulcus – which is the area of developing heart first being covered by the epicardium (Vrancken

Peeters *et al.*, 1997b; Kattan *et al.*, 2004).

The origin of the erythrocytes within the blood islands is another interesting issue. It has been suggested that the erythrocytes and angioblasts derive from a common precursor cells – hemangioblasts (Pardanaud *et al.*, 1989; Muñoz-Chápuli *et al.*, 1999). Retroviral tagging of the proepicardium showed that the erythrocytes of the subepicardium derive from the proepicardial progenitor cells which differentiate into erythroblasts within the heart (Tomanek *et al.*, 2006a). This appears to be consistent with the results of earlier studies showing the presence of erythroblasts within the subepicardium (Virágh 1990; Kálmán *et al.*, 1995) and the recent study by Kattan (2004) demonstrating presence of progenitor cell markers (CD45) on erythroblast clusters within subepicardial area. Another possible source of erythrocytes within the blood islands, would be the systemic circulation from where they migrate to the heart and subsequently assemble with angioblasts (Ratajska *et al.*, 2006).

Retroviral tagging of the proepicardium identified the sources of smooth muscle cells of the tunica media, fibroblasts of the adventitia of large vessels, and the interstitial heart fibroblasts (Mikawa and Fischman, 1992; Mikawa and Gourdie, 1996). All these cell types derive from the epicardial mesothelium by the epithelial-mesenchymal transformation (EMT) and subsequent differentiation (Dettman *et al.*, 1998; Gittenberger-de Groot *et al.*, 1998; Vrancken Peeters *et al.*, 1999; Pérez-Pomares *et al.*, 2002a; Muñoz-Chápuli *et al.*, 2002). Mesenchymal cells are able to infiltrate myocardium reaching the subendocardial cushions (Gittenberger-de Groot *et al.*, 1998). Depending on local availability of the growth factors such as PDGF-BB, VEGF or bFGF, the mesenchymal cells differentiate into smooth muscle cells or endothelial cells, respectively (Cox, Poole 2000; Pérez-Pomares *et al.*, 2002a).

The studies of chicken-quail chimeras broadened our understanding of the precursors and development of the coronary vessel. There are two main theories on the origin of the vascular endothelial cells. First theory, based on experiments with quail-chicken chimeras in which small pieces of quail liver were transplanted into the chicken coelom, claims that vascular endothelial cells derive from liver primordium, and the angioblasts differentiate within the subepicardium which subsequently gives rise to the primitive subepicardial vascular plexus (Poelmann *et al.*, 2002). The second theory maintains that epicardium-derived mesenchymal cells (EPDC) are able to differentiate into vascular endothelial cells (Männer 1999; Muñoz-Chápuli *et al.*, 2002; Pérez-Pomares *et al.*, 2002a). The latter theory is supported by the results of the retroviral labeling of the proepicardium, and by the chicken-quail




Fig. 5. The epicardium of a mouse heart at stage 13 dpc. Vascular vesicles located in the subepicardial space contain nucleated red blood cells (arrows). (A) Section of the ventricles and the interventricular sulcus; the area in the rectangle is enlarged in (B), mag. 150x. (C) Vascular vesicle in the interventricular sulcus. These vesicles are precursors of subepicardial coronary vessels. Mag. 380x in (B,C).

chimera studies. Double labeling of EPDC with anti-QH1 antibodies and cytokeratin showed that both markers were present in the precursors of the coronary endothelial cells. Additional support for this theory comes from the interspecies comparison of epicardially derived cells. In dogfish (*Scyliorhinus canicula*) all subepicardially derived cells originate from the epicardial mesothelium. In this species subepicardial vessels arise long before the migration of extracardiac angioblasts begins (Muñoz-Chápuli *et al.*, 1996). The experiments in rats in which avascular hearts were grafted *in oculo* showed that the coronary vessels in transplanted hearts develop without any vascular contribution from the host (Rongish *et al.*, 1994). Both studies are in agreement with the idea that splanchnic angioblast differentiation occurs *in situ* without significant migration of vascular cell precursors (Drake *et al.*, 1997). The idea that the angioblasts originate *in situ* from the splanchnic mesoderm is also supported by the studies of 5-somite stage avian embryo (Cox and Poole 2000; Poole *et al.*, 2001).

As we mentioned previously, the blood islands are precursors of coronary vasculature. Blood islands differentiate into vascular tubes (capillaries), which appear first within the subepicardium. Capillaries grow within the subepicardium of the atrioventricular and interventricular sulcuses and coalesce forming primitive vascular network. Formation of the coronary arteries commences with directional growth of the protruding capillaries towards the sinuses of Valsalva and establishing patency with the aortic lumen (Bogers *et al.*, 1990; Waldo *et al.*, 1990). The protrusion of primordia of coronary arteries towards the aorta is facilitated by apoptosis of the aortic wall (Bernanke, Valkey 2002; Tomanek *et al.*, 2006b). Connection with the aorta is established in rats on ED16 (Heinzberger 1983), at HH stage 29-35 in chicken and quail hearts (Kattan *et al.*, 2004), and at 44-49 day in human heart (Conte, Pellegrini 1984; Hutchins *et al.*, 1988; Mandarin-Lacerda 1990). Interestingly, there are several capillaries penetrating towards the three sinuses of Valsalva but only two (in the left and right sinuses, respectively) establish connection with the aorta (Poelmann *et al.*, 1993). Capillary plexus spreads around the outflow tract and within the myocardium of diaphragmatic surface of the heart. Subsequently, the plexus grows on the sternal surface of the heart. The left ventricle undergoes vascularization followed by the vascularization of the right ventricle. The apex of the heart vascularizes as the last (Rychter *et al.*, 1975; Vrancken-Peeters *et al.*, 1997b, Kattan *et al.*, 2004). Initially there is a gradient of the myocardial wall vascularization with the highest density of vessels in the subepicardium and the lowest in the subendocardium. This is consistent with the expression of VEGF and its receptor-2 (Tomanek *et al.*, 1999a; 2002; Tomanek, Zheng 2002). It is believed that the thickening of the myocardial wall (Tomanek *et al.*, 1999b), and relative hypoxia in the subepicardium are the stimuli responsible for the rapid development of coronary vessels in the subepicardium. Multiple growth factors participate in the development and spreading of the vasculature (Ratajska *et al.*, 1995; Tomanek *et al.*, 1998; 2001). We do not know what signal allows the directional growth of coronary artery precursors towards the aorta. Probably the aorta is recognized by at least one signal from growth factors or from parasympathetic ganglia situated at the roots of the aorta (Hood, Rosenquist 1992). The parasympathetic ganglia which are derivative of neural crest cells (Verberne *et al.*, 1998) are positioned closely to coronary artery ostia in chick heart (Waldo *et al.*, 1994). This suggests that the




Fig. 6. Rat heart of 18 dpc embryo. Immunostaining with anti-smooth muscle myosin antibody (A,B) smooth muscle myosin is present within the proximal (A) and distal wall (B) of coronary artery.

signal(s) from the parasympathetic ganglia may contribute to the formation of coronary ostia. In cases such as Bland-White-Garland syndrome, this signal is imprecise allowing the primordia to grow towards the pulmonary trunk (1933). In congenital heart malformations, however, with dextroposition of the aorta relative to the pulmonary trunk, the primordial coronary arteries grow towards the aorta (González-Iriarte *et al.*, 2003; Ratajska *et al.*, 2005).

Formation of the tunica media of larger coronary vessels starts at the proximal end of coronary artery by building up a layer of mesenchymal cells. These cells approach the vascular wall before the connection of coronary artery precursors with the aorta is established (Ratajska, Fiejka 1999). Further differentiation into smooth muscle cells commences when blood starts to flow within the patent coronary system (Hood, Rosenquist 1992). The differentiation into smooth muscle cells requires acquisition of certain contractile and cytoskeletal proteins, which occurs sequentially and is species-specific (Owens *et al.*, 1995; Hungerford, Little 1999; Ratajska *et al.*, 2001). The first is the smooth muscle alpha actin, which appears sequentially from the proximal to distal part of the coronary vessel wall (Hood, Rosenquist 1992, Ratajska *et al.*, 2001). Subsequently, the smooth muscle myosin appears

(Fig. 6). Concomitantly with the start of blood flow within the patent coronary system, the shear stress on endothelial cells increases stimulating a release of certain substances from the epithelial cells (Hannan *et al.*, 1988; Joseph-Silverstein *et al.*, 1987). These substances modulate smooth muscle cell differentiation and influence the smooth muscle cell proliferation (Sterpetti *et al.*, 1993). An increase in shear stress within the arterial side of the coronary system and rapid blood flow in arterial wall explains why the vascular wall of arterial system differentiates earlier than the cardiac veins (Vrancken Peeters *et al.*, 1997a). In addition, the specific shape of the internal wall of coronary vasculature of the embryonic heart can be explained by the elevated speed of blood flow within the arteries as compared to veins (Ratajska *et al.*, 2003). The development of the coronary vessel is not restricted to the embryonic period of life and it continues postnatally. It is accomplished by remodeling of the vasculature: spreading of existing vessels by sprouting, intussusceptive growth and disappearing of the branches connecting the vessels (Tomanek 2005).

The role of the epicardium in myocardial development

Formation of the epicardium is necessary for normal development of myocardium, its "tightness", stiffness, elasticity, and for the proper vascularization of the heart wall. The heart devoid of the proepicardium (as the result of photoablation) develops malformations such as double outlet right ventricle, and deficiency of the interventricular septum in its membranous part (Männer *et al.*, 2005). It is highly probable that EPDC takes part in the formation of the interventricular septum. It is not known however, in what way EPDC could induce the complex pathological changes of the heart such as double outlet right ventricle (Männer *et al.*, 2005). Mice devoid of *RXRalpha* gene exhibit delayed development of the epicardium. Their epicardium exhibits malformations with exuberant extracellular matrix and increased apoptosis of mesenchymal cells (Jenkins *et al.*, 2005). These mice have lower number of subepicardial mesenchymal cells. In addition, the fibronectin aggregation in their subepicardium is abnormal (Jenkins *et al.*, 2005).

The mesenchymal cells (deriving from the epicardial cells) are able to migrate from the epicardium to the subendocardial area where they differentiate into fibroblasts, which synthesize extracellular matrix. These mesenchymal cells migrate to the subendocardial area where they form heart valves. They also play an active role in the formation of the fibrous heart skeleton by inducing cardiomyocyte-fibroblast transformation, which occurs in areas of insulation of atrial and ventricular myocardium (Winter, Gittenberger-de Groot 2007).

The recent studies on chicken embryos, indicate an inductive influence of EPDC on those cardiac myocytes that differentiate into the conductive system of the heart (Gittenberger-de Groot *et al.*, 2003) and the periarterial and subendocardial Purkinje fibers (Gourdie *et al.*, 1998, 1999). EPDC are also able to differentiate into smooth muscle cells of the tunica media and fibroblasts of the adventitia (Vrancken Peeters *et al.*, 1999). This suggests that the inductive influence of EPDC on Purkinje fiber differentiation is indirect and does not require a direct contact with the arteries and periarterial Purkinje fibers. *In vitro* studies demonstrated that endothelin cascade induces Purkinje fiber differentiation (Gourdie *et al.*, 1998). This has been confirmed by *in vivo* studies

(Takebayashi-Suzuki *et al.*, 2000), which showed that the communication between EPDC and primitive cardiocytes is mediated via endothelins released from EPDC. However, in birds, the EPDC play a more direct role in differentiation of periarterial and subendocardial Purkinje fibers of the heart.

In some rare cases the epicardial mesothelium undergoes a malignant transformation into a tumor – mesothelioma. It is a very rare tumor, which derives from membranous serosa of body cavities, and rarely from pericardium or from the epicardium (Hammar 2006). In the latter case, the tumor grows into the pericardial cavity and surrounds the heart. Microscopically, this tumor consists of epithelial cells, which sometimes form canalicular, glandular or papillomatous structures.

Acknowledgments

This paper is supported by Ministry of Education and Science grant (#2P05A 111 28) and internal funds of the Medical University of Warsaw. The technical help of A. Podbielska and M. Michniewska is much appreciated.

References

- BERNANKE, D.H., VELKEY, J.M. (2002) Development of the coronary blood supply: changing concepts and current ideas. *Anat Rec (New Anat)* 269: 198-208.
- BLAND, E.F., WHITE, P.D., and GARLAND, J. (1933) Congenital anomalies of coronary arteries: report of an unusual case associated with cardiac hypertrophy. *Am Heart J* 8: 787-801.
- BOGERS, A.J.J.C., GITTENBERGER DE GROOT, A.C., POELMANN, R.E., PEULT, B.M. and HUYSMANS, H.A. (1989) Development of the origin of the coronary arteries, a matter of ingrowth or outgrowth? *Anat Embryol* 180: 437-441.
- CANO, A., PÉREZ-MORENO, M., RODRIGO, I., LOCASCIO, A., BLANCO, M.J., DEL BARRIO, M.G., PORTILLO, F. and NIETO, M.A. (2000) The transcription factor snail controls epithelial-mesenchymal transitions by repressing E-cadherin expression. *Nature Cell Biol* 2: 76-83.
- CARMONA, R., GONZÁLEZ-IRIARTE, M., MACÍAS, D., PÉREZ-POMARES, J.M., GARCÍA-GARRIDO, L., MUÑOZ-CHÁPULI, R. (2000) Immunolocalization of the transcription factor Slug in the developing avian heart. *Anat Embryol* 201: 103-109.
- CONTE, G., PELLEGRINI, A. (1984) On the development of the coronary arteries in human embryos, stages 14-19. *Anat Embryol* 169: 209-218.
- COX, C.M., POOLE, T.J. (2000) Angioblasts differentiation is influenced by the local environment: FGF-2 induces angioblasts and patterns vessel formation in the quail embryo. *Dev Dyn* 218: 371-382.
- DETTMAN, R.W., DENETCLAW, W., ORDAHL, C.P. and BRISTOW, J. (1998) Common epicardial origin of coronary vascular smooth muscle, perivascular fibroblasts, and intermyocardial fibroblasts in the avian heart. *Dev Biol* 193: 169-181.
- DRAKE, C.J., BRANDT, S.J., TRUSK, T.C. and LITTLE, C.D. (1997) TAL1/SCL is expressed in endothelial progenitor cells/angioblasts and defines a dorsal-ventral gradient of vasculogenesis. *Dev Biol* 192: 17-30.
- GITTENBERGER-DE GROOT, A.C., VRANCKEN PEETERS, M.P.F.M., MENTINK, M.M.T., GOURDIE, R.G. and POELMANN, R.E. (1998) Epicardium-derived cells contribute a novel population to the myocardial wall and the atrioventricular cushions. *Circ Res* 82: 1043-1052.
- GITTENBERGER-DE GROOT, A.C., BLOM, N.M., AOYAMA, N., SUCOV, H., WENINK, A.C.G. and POELMANN, R.E. (2003) The role of neural crest and epicardium-derived cells in conduction system formation. *Novartis Foundation Symposium* 250: 125-141.
- GONZÁLEZ-IRIARTE, M., CARMONA, R., PÉREZ-POMARES, J.M., MACÍAS, D., COSTELL, M. and MUÑOZ-CHÁPULI, R. (2003) Development of the coronary arteries in murine model of transposition of the great arteries. *J Mol Cell Cardiol* 35: 795-802.

- GOURDIE, R.G., WEI, Y., KIM, D., KLATT, S.C. and MIKAWA, T. (1998) Endothelin-induced conversion of embryonic heart muscle cells into impulse-conducting Purkinje fibers. *Proc Natl Acad Sci USA* 95: 6815-6818.
- GOURDIE, R.G., KUBALAK, S.W. and MIKAWA, T. (1999) Conduction the embryonic heart: orchestrating development of specialized cardiac tissue. *Trends Cardiovasc Med* 9: 18-26.
- GUADIX, J., CARMONA, R., MUÑOZ-CHÁPULI, R. and PÉREZ-POMARES, J.M. (2006) In vivo and in vitro analysis of the vasculogenic potential of avian proepicardial and epicardial cells. *Dev Dyn* 235: 1014-1026.
- HAMBURGER, V., HAMILTON, H.L. (1951) A series of normal stages in the development of the chick embryos. *J Morphol* 88: 49-92.
- HAMMAR S.P. (2006) Macroscopic, histologic, histochemical, immunohistochemical, and ultrastructural features of mesothelioma. *Ultrastruct Pathol* 30: 3-17.
- HANNAN, R.L., KOUREMBANAS, S., FLANDERS, K.C., ROGELJ, S.J., ROBERTS, A.B., FALLER, D.V. and KLAGSBRUN, M. (1988) Endothelial cells synthesize basic fibroblast growth factor and transforming growth factor beta. *Growth Factors* 1: 7-17.
- HATCHER, C.J., DIMAN, NYS-G, KIM, M.-S., PENNISI, D., SONG, Y., GOLDSTEIN, M.M., MIKAWA, T. and BASSON, C.T. (2004) A role for Tbx5 in proepicardial cell migration during cardiogenesis. *Physiol Genomics* 18: 129-140.
- HEINZBERGER, C.F.M. (1983) Development of myocardial vascularization in the rat. *Acta Morphol Neerl-Scand* 21: 267-284.
- HIRAKOW, R. (1983) Development of the cardiac blood vessels in staged human embryos. *Acta Anat* 115: 220-230.
- HIRUMA, T., HIRAKOW, R. (1989) Epicardial formation in embryonic chick heart: computer-aided reconstruction, scanning, and transmission electron microscopic studies. *Am J Anat* 184: 129-138.
- HOOD, L.C., ROSENQUIST, T.H. (1992) Coronary artery development in the chick: origin and deployment of smooth muscle cells, and the effects of neural cell ablation. *Anat Rec* 234: 291-300.
- HUNGERFORD, J.E., LITTLE, C.D. (1999) Developmental biology of the vascular smooth muscle cell: building a multilayered vessel wall. *J Vasc Res* 36:2-27.
- HUTCHINS, G.M., KESSLER-HANNA, A. and MOORE, G.W. (1988) Development of the coronary arteries in the embryonic human heart. *Circulation* 77: 1250-1257.
- HO, E., SHIMADA, Y. (1978) Formation of the epicardium studied with the scanning electron microscope. *Dev Biol* 66: 579-585.
- JENKINS, S.J., HUTSON, D.R. and KUBALAK, S.W. (2005) Analysis of the proepicardium-epicardium transition during the malformation of the RXRalpha-epicardium. *Dev Dyn* 233: 1091-1101.
- JOSEPH-SILVERSTEIN, J., RIFKIN, D.B. (1987) Endothelial cell growth factors and the vessel wall. *Semin Thromb Hemostasis* 13: 504-513.
- KÁLMÁN, F., VIRÁGH, S. and MODIS, L. (1995) Cell surface glycoconjugates and the extracellular matrix of the developing mouse embryo epicardium. *Anat Embryol* 191: 451-464.
- KATTAN, J., DETTMAN, R.W. and BRISTOW, J. (2004) Formation and remodeling of the coronary vascular bed in the embryonic avian heart. *Dev Dyn* 230: 34-43.
- KUHN, H.-J., LIEBHERR, G. (1988) The early development of the epicardium in *Tupaia belangeri*. *Anat Embryol* 177: 225-234.
- KURKIEWICZ, T. (1909) O histogenezie mieŚnia sercowego zwierzat kregowych. *Bull Int Acad Sci Cracovie*, 148-191.
- KWEE, L., BALDWIN, H.S., SHEN, H.M., STEWART, C.L., BUCK, C., BUCK, C.A. and LABOW, M.A. (1995) Defective development of the embryonic and extraembryonic circulatory systems in vascular cell adhesion molecule (VCAM-1) - deficient mice. *Development* 121: 489-503.
- MARKWALD, R.R., FITZHARRIS, T.P. and MANASEK, F.J. (1977) Structural development of endocardial cushions. *Am J Anatomy* 148: 85-119.
- MANDARIM-DE-LACERDA, C.A. (1990) Development of the coronary arteries in staged human embryos (The Paris Embryological Collection Revisited) *Ann Acad Bras Ci* 62:79-84.
- MÄNNER, J. (1992) The development of the pericardial villi in the chick embryo. *Anat Embryol* 186: 379-385.
- MÄNNER, J. (1993) Experimental study on the formation of the epicardium in chick embryos. *Anat Embryol* 187: 281-289.
- MÄNNER, J. (1999) Does the subepicardial mesenchyme contribute myocardioblasts to the chick embryo heart? A quail-chick chimera study tracing the fate of the epicardial primordium. *Anat Rec* 255: 212-226.
- MÄNNER, J., PÉREZ-POMARES, J.M., MACÍAS, S.D. and MUÑOZ-CHÁPULI, R. (2001) The origin, formation and developmental significance of the epicardium: a review. *Cell Tissue Organs* 169: 89-103.
- MÄNNER, J., SCHLUETER, J., BRAND, T. (2005) Experimental analyses of the function of the proepicardium using a new microsurgical procedure to induce loss-of-proepicardial-function in chick embryos. *Dev Dyn* 233: 1454-1463.
- MIKAWA, T., FISCHMAN, D.A. (1992) Retroviral analysis of cardiac morphogenesis: discontinuous formation of coronary vessels. *Proc Natl Acad Sci USA* 89: 9504-9508.
- MIKAWA, T., GOURDIE, R.G. (1996) Pericardial mesoderm generates a population of coronary smooth muscle cells migrating into the heart along with ingrowth of the epicardial organ. *Dev Biol* 174: 221-232.
- MOORE, A.W., MCINNES, L., KREIDBERG, J., HASTIE, N.D. and SCHEDL, A. (1999) YAC complementation shows a requirement for WT1 in development of epicardium, adrenal gland and throughout nephrogenesis. *Development* 126: 1845-1857.
- MUÑOZ-CHÁPULI, R., GONZÁLEZ-IRIARTE, M., CARMONA, R., ATENCIA, G., MACÍAS, D. and PÉREZ-POMARES, J.M. (2002) Cellular precursors of the coronary arteries. *Tex Heart Inst J* 29: 243-249.
- MUÑOZ-CHÁPULI, R., MACÍAS, D., RAMOS, C., GALLEGO, A. and ANDRÉS, V. (1996) Development of the subepicardial mesenchyme and the early cardiac vessels in the dogfish (*Scyliorhinus canicula*). *J Exp Zool* 275: 95-111.
- MUÑOZ-CHÁPULI, R., PÉREZ-POMARES, J.M., MACÍAS, D., GARCÍA-GARRIDO, L., CARMONA, R. and GONZÁLEZ, M. (1999) Differentiation of hemangioblasts from embryonic mesothelial cells? A model of the origin of the vertebrate cardiovascular system. *Differentiation* 64: 133-141.
- NAHIRNEY, P.C., MIKAWA, T. and FISCHMAN, D.A. (2003) Evidence for an extracellular matrix bridge guiding proepicardial cell migration to the myocardium of chick embryos. *Dev Dyn* 227: 511-523.
- NESBITT, T., LEMLEY, A., DAVIS, J., YOST, M.J., GOODWIN, R.L. and POTTS, J.D. (2006) Epicardial development in the rat: a new perspective. *Microsc Microanal* 12: 390-398.
- OLIVEY, H.E., MUNDELL, N.A., AUSTIN, A.F. and BARNETT, J.V. (2006) Transforming growth factor-beta stimulates epithelial-mesenchymal transformation in the proepicardium. *Dev Dyn* 235: 50-59.
- OWENS, G.K. (1995) Regulation of differentiation of vascular smooth muscle cells. *Physiol Rev* 75: 487-517.
- PARDANAUD, L., YASSINE, F. and DIETERLEN-LIEVRE, F. (1989) Relationship between vasculogenesis, angiogenesis and haemopoiesis during avian ontogeny. *Development* 105: 473-485.
- PÉREZ-POMARES, J.M., MACÍAS, D., GARCÍA-GARRIDO, L. and MUÑOZ-CHÁPULI, R. (1998) The origin of the subepicardial mesenchyme in the avian embryo: an immunohistochemical and quail-chimera study. *Dev Biol* 200: 57-68.
- PÉREZ-POMARES, J.M., CARMONA, R., GONZALEZ-IRIARTE, M., ATENCIA, G., WESSELS, A. and MUÑOZ-CHÁPULI, R. (2002a) Origin of coronary endothelial cells from epicardial mesothelium in avian embryos. *Int J Dev Biol* 46: 1005-1013.
- PÉREZ-POMARES, J.M., MIRONOV, V., GUADIX, J.A., MACÍAS, D., MARKWALD, R.R., MUÑOZ-CHÁPULI, R. (2006) In vitro self-assembly of proepicardial cell aggregates: an embryonic vasculogenic model for vascular tissue engineering. *Anat Rec Part A* 288A: 700-713.
- PÉREZ-POMARES, J.M., PHELPS, A., SEDMEROVA, M., CARMONA, R., GONZALEZ-IRIARTE, M., MUÑOZ-CHÁPULI, R. and WESSELS, A. (2002b) Experimental studies on the spatiotemporal expression of WT1 and RALDH2 in the embryonic avian heart: a model for the regulation of myocardial and valvuloseptal development by epicardially derived cells (EPDCs). *Dev Biol* 247: 307-326.
- POELMANN, R.E., GITTEBERGER-DE GROOT, A.C., MENTINK, M.M.T., BOKENKAMP, R. and HOGERS, B. (1993) Development of the cardiac coronary vesicular endothelium, studied with antiendothelial antibodies, in chicken-quail chimeras. *Circ Res* 73: 559-568.
- POELMANN, R.E., LIE-VENEMA, H., GITTEBERGER-DE GROOT, A.C. (2002)

- The role of the epicardium and neural crest as extracardiac contributors to coronary vascular development. *Texas Heart Inst J*29: 255-261.
- POOLE, T.J., FINKELSTEIN, E.B. and COX, C.M. (2001) The role of FGF and VEGF in angioblasts induction and migration during vascular development. *Dev Dyn* 220: 1-17.
- RATAJSKA, A., CISZEK, B. and SOWINSKA, A. (2003) Embryonic development of coronary vasculature in rats: corrosion casting studies. *Anat Rec* 270A(2): 109-116.
- RATAJSKA, A., FIEJKA, E. (1999) Prenatal development of coronary arteries in the rat: morphologic pattern. *Anat Embryol* 200: 533-540.
- RATAJSKA, A., CZARNOWSKA, E., KOLODZINSKA, A., KLUZEK, W. and LESNIAK, W. (2006) Vasculogenesis of the embryonic heart: origin of blood-island-like structures. *Anat Rec A* 288: 223-232.
- RATAJSKA, A., TORRY, R.T., KITTEN, G.T., KOLKER, S.J. and TOMANEK, R.J. (1995) Modulation of cell migration and vessel formation by vascular endothelial growth factor and basic fibroblast growth factor in cultured embryonic heart. *Dev Dyn* 203: 399-407.
- RATAJSKA, A., ZLOTOROWICZ, R., BLAZEJCZYK, M. and WASIUTYNSKI, A. (2005) Coronary artery embryogenesis in cardiac defects induced by retinoic acid in mice. *Birth Defects Res A Clin Mol Teratol* 73: 966-979.
- RATAJSKA, A., ZARSKA, M., QUENSEL, C. and KRÄMER, J. (2001) Differentiation of the smooth muscle cell phenotypes during embryonic development of coronary vessels in the rat. *Histochem Cell Biol* 116: 79-87.
- RONGISH, B.J., TORRY, R.J., TUCKER, D.C. and TOMANEK, R.J. (1994) Neovascularization of embryonic rat hearts cultured *in oculo* closely mimics *in utero* coronary vessel development. *J Vasc Res* 31: 205-21.
- RYCHTER, Z., JIRÁSEK, J.E., RYCHTEROVÁ, V. and UHER, J. (1975) Vascularization of heart in human embryo: location and shape of non-vascularized part of cardiac wall. *Folia Morphol (Prague)* 23: 88-96.
- SCHLUETER, J., MÄNNER, J. and BRAND, T. (2006) BMP is important regulator of proepicardial identity in the chick embryo. *Dev Biol* 295: 546-558.
- SCHULTE, I., SCHLUETER, J., ABU-ISSA, R., BRAND, T. and MÄNNER, J. (2007) Morphological and molecular left-right asymmetries in the development of the proepicardium: a comparative analysis on mouse and chick embryos. *Dev Dyn* 236: 684-95.
- STERPETTI, A.V., CUCINA, A., D'ANGELO, L.S., CARDILLO, B., CAVALLARO, A. (1993) Shear stress modulates the proliferation rate, protein synthesis, and mitogenic activity of arterial smooth muscle cells. *Surgery* 113: 691-699.
- TAKEBAYASHI-SUZUKI, K., YANAGISAWA, M., GOURDIE, R.G., KANZAWA, N. and MIKAWA, T. (2000) In vivo induction of cardiac Purkinje fiber differentiation by coexpression of preproendothelin-1 and endothelin converting enzyme-1. *Development* 127: 3523-3532.
- TOMANEK, R.J. (2005) Formation of coronary vasculature during development. *Angiogenesis* 8: 273-284.
- TOMANEK, R.J., ISHII, Y., HOLIFIELD, J.S., SJOGREN, C.L., HANSEN, H.K. and MIKAWA, T. (2006a) VEGF family members regulate myocardial tubulogenesis and coronary artery formation in the embryo. *Circ Res* 98: 947-953.
- TOMANEK, R.J., HANSEN, H.K. and DEDKOV, E.I. (2006b) Vascular patterning of the quail coronary system during development. *Anat Rec* 288A: 989-999.
- TOMANEK, R.J., HU, N., PHAN, B. and CLARK, E.D. (1999b) Rate of coronary vascularization during embryonic chicken development is influenced by the rate of myocardial growth. *Cardiovasc Res* 41: 663-671.
- TOMANEK, R.J., HOLIFIELD, J.S., REITER, R.S., SANDRA, A. and LIN, J.J. (2002) Role of VEGF family members and receptors in coronary vessel formation. *Dev Dyn* 225:233-240.
- TOMANEK, R.J., LOTUN, K., CLARK, E.B., SUVARNA, P.R., HU, N. (1998) VEGF and bFGF stimulate myocardial vascularization in embryonic chick. *Am J Physiol* 274: H1620-1626.
- TOMANEK, R.J., RATAJSKA, A., KITTEN, G.T., YUE, X. and SANDRA, A. (1999a) Vascular endothelial growth factor expression coincides with coronary vasculogenesis and angiogenesis. *Dev Dyn* 215: 54-61.
- TOMANEK, R.J., ZHENG, W. (2002) Role of growth factors in coronary morphogenesis. *Tex Heart Inst J*29:250-254.
- TOMANEK, R.J., ZHENG, W., PETERS, K.G., LIN, P., HOLIFIELD, J.S. and SUVARNA, P.R. (2001) Multiple growth factors regulate coronary embryonic vasculogenesis. *Dev Dyn* 221: 265-273.
- VAN DEN EIJNDE, S.M., WENINK, A.C., VERMEIJ-KEERS, C. (1995) Origin of subepicardial cells in rat embryos. *Anat Rec* 242: 96-102.
- VIRÁGH, S., CHALLICE, C.E. (1981) The origin of the epicardium and the embryonic myocardial circulation in the mouse. *Anat Rec* 201: 157-168.
- VIRÁGH, S., GITTENBERGER-DE GROOT, A., POELMANN, R.E. and KÁLMÁN, F. (1993) Early development of quail heart epicardium and associated vascular and glandular structures. *Anat Embryol* 188: 381-393.
- VIRÁGH, S., KÁLMÁN, F., GITTENBERGER-DE GROOT, A.C., POELMANN, R.E. and MOORMAN, A.F.M. (1990) Angiogenesis and hematopoiesis in the epicardium of the vertebrate embryo heart, w: "Embryonic origins of defective heart development", Bockman DE, Kirby ML (wyd). *Ann NY Acad Sci USA* 588: 455-458.
- VERBERNE, M.E., GITTENBERGER-DE GROOT, A.C., POELMANN, R.E. (1998) Lineage and development of the parasympathetic nervous system of the embryonic chick heart. *Anat Embryol* 198: 171-184.
- VRANCKEN PEETERS, M.P.F.M., MENTINK, M.M.T., POELMANN, R.E., GITTENBERGER-DE GROOT, A.C. (1995) Cytokeratins as a marker for epicardial formation in the quail embryo. *Anat Embryol* 191: 503-508.
- VRANCKEN PEETERS, M.P.F.M., GITTENBERGER-DE GROOT, A.C., MENTINK, M.M.T., HUNGERFORD, J.E., LITTLE, C.D. and POELMANN, R.E. (1997a) Differences in development of coronary arteries and veins. *Cardiovasc Res* 36: 101-110.
- VRANCKEN PEETERS, M.P.F.M., GITTENBERGER-DE GROOT, A.C., MENTINK, M.M.T., HUNGERFORD, J.E., LITTLE, C.D. and POELMANN, R.E. (1997b) The development of the coronary vessels and their differentiation into arteries and veins in the embryonic quail heart. *Dev Dyn* 208: 338-348.
- VRANCKEN PEETERS, M.P.F.M., GITTENBERGER-DE GROOT, A.C., MENTINK, M.M.P. and POELMANN, R.E. (1999) Smooth muscle cells and fibroblasts of the coronary arteries derive from epithelial-mesenchymal transformation of the epicardium. *Anat Embryol* 199: 367-378.
- WALDO, K.L., KUMINSKI, D.H., KIRBY, M.L. (1994) Association of the cardiac neural crest with development of the coronary arteries in the chick embryo. *Anat Rec* 239: 315-331.
- WALDO, K.L., WILLNER, W. and KIRBY, M.L. (1990) Origin of the proximal coronary artery stems and a review of ventricular vascularization in the chick embryo. *Am J Anat* 188: 109-120.
- WATT, A.J., BATTLE, M.A., DUNCAN, S.A. (2004) GATA4 is essential for formation of the proepicardium and regulates cardiogenesis. *Proc Natl Acad Sci USA* 101: 12573-12578.
- WESSELS, A., PÉREZ-POMARES, J.M. (2004) The epicardium and epicardially derived cells (EPDCs) as cardiac stem cells. *Anat Rec* 276:43-57.
- WINTER, E.M., GITTENBERGER-DE GROOT, A.C. (2007) Epicardium-derived cells in cardiogenesis and cardiac regeneration. *Cell Mol Life Sci* 64: 692-703.

Related, previously published *Int. J. Dev. Biol.* articles

See our recent Special Issue ***Developmental Biology in Poland*** edited by Kloc, Maleszewski and Tarkowski at: <http://www.ijdb.edu.es/web/contents.php?vol=52&issue=2-3>

See our Special Issue ***Mammalian Reproduction & Development*** in honor of Anne McLaren and edited by Brigid Hogan at: <http://www.ijdb.edu.es/web/contents.php?vol=45&issue=3>

An efficient method for isolation of murine bone marrow mesenchymal stem cells

Samad Nadri, Masoud Soleimani, Reza H. HosSeni, Mohammad Massumi, Amir Atashi and Reza Izadpanah
Int. J. Dev. Biol. (2007) 51: 723-729

Enhanced development of porcine embryos cloned from bone marrow mesenchymal stem cells

Hai-Feng Jin, B. Mohana Kumar, Jung-Gon Kim, Hye-Jin Song, Yeon-Ji Jeong, Seong-Keun Cho, Sivasankaran Balasubramanian, Sang-Yong Choe and Gyu-Jin Rho
Int. J. Dev. Biol. (2007) 51: 85-90

Interleukin-2 induces the proliferation of mouse primordial germ cells in vitro

Cristina Eguizabal, Maria D. Boyano, Alejandro Díez-Torre, Ricardo Andrade, Noelia Andollo, Massimo De Felici and Juan Aréchaga
Int. J. Dev. Biol. (2007) 51: 731-738

Tracing the hemangioblast during embryogenesis: developmental relationships between endothelial and hematopoietic cells

Thierry Jaffredo, Karine Bollerot, Daisuke Sugiyama, Rodolphe Gautier and Cécile Drevon
Int. J. Dev. Biol. (2005) 49: 269-277

Of birds and mice: hematopoietic stem cell development

Isabelle Godin and Ana Cumano
Int. J. Dev. Biol. (2005) 49: 251-257

Embryonic development of the human hematopoietic system

Manuela Tavian and Bruno Péault
Int. J. Dev. Biol. (2005) 49: 243-250

Vasculogenesis and angiogenesis in the mouse embryo studied using quail/mouse chimeras

Michel Pudliszewski and Luc Pardanaud
Int. J. Dev. Biol. (2005) 49: 355-361

Developmental expression of Pod 1 in *Xenopus laevis*

Subreena Simrick, Karine Massé and Elizabeth A. Jones
Int. J. Dev. Biol. (2005) 49: 59-63

Origin of coronary endothelial cells from epicardial mesothelium in avian embryos.

José-María Pérez-Pomares, Rita Carmona, Mauricio González-Iriarte, Gerardo Atencia, Andy Wessels and Ramón Muñoz-Chápuli
Int. J. Dev. Biol. (2002) 46: 1005-1013

The contribution of the proepicardium to avian cardiovascular development

JM Perez-Pomares, A Phelps, R Munoz-Chapuli, A Wessels
Int. J. Dev. Biol. (2001) 45: S155-S156

Origin of endothelial cells from mesothelial-derived mesenchymal cells in the liver of avian embryos

R Munoz-Chapuli, R Carmona, M Gonzalez-Iriarte, JM Perez-Pomares, D Macias, G Atencia, MJ Aranda
Int. J. Dev. Biol. (2001) 45: S143-S144



2006 ISI **Impact Factor = 3.577**

