

# Comparative epigenetic evaluation of human embryonic stem and induced pluripotent cells

RAHA FAVAEDI<sup>1</sup>, MARYAM SHAHHOSEINI<sup>\*1</sup>, MAHAMMAD PAKZAD<sup>2</sup>, SEPIDEH MOLLAMOHAMMADI<sup>2</sup>  
and HOSSEIN BAHARVAND<sup>2</sup>

<sup>1</sup>Department of Genetics, Reproductive Biomedicine Research Center, Royan Institute for Reproductive Biomedicine, ACECR, Tehran, Iran and <sup>2</sup>Department of Stem Cells and Developmental Biology, Cell Science Research Center, Royan Institute for Stem Cell Biology and Technology, ACECR, Tehran, Iran.

**ABSTRACT** Histone H3 lysine 9 methylation has been shown to be a critical barrier to efficient cell reprogramming. This discovery allows the assessment of the cell pluripotency state by considering the extent of H3K9 methylation vs. acetylation at the same position. A set of pluripotent and differentiated human cells including embryonic stem cells, their differentiated and reprogrammed counterparts, along with human fibroblasts and their derived reprogrammed cells, were used to evaluate the ratio of total H3K9 methylation over acetylation using a quantitative ELISA-based approach. Also, the occurrence of the H3K4me3 and H3K27me3 bivalent marks was evaluated. Additionally, using ChIP-qPCR the occurrence of these histone marks on the regulatory regions of stemness genes (*Nanog*, *Oct4* and *Sox2*) as well as on genes indicating fibroblast differentiation (*Vim*, *COL1A1* and *THY1*) was evaluated. We evidence remarkably high ratios of H3K9ac/K9me2 in ES and iPSC cells vs. differentiated cells. In iPSCs, a direct relationship between the ratios of total H3K9ac/H3K9me2 and the ratios of these marks on pluripotency gene regulatory regions and their expression was observed. In differentiated cells, in contrast, the ratios of global H3K9ac/K9me2 is low but the active genes escape this general situation and bear higher amounts of H3K9ac vs. H3K9me. Total H3K4me3/K27me3 ratios presented the same trends, but with reduced amplitudes. We propose that the rapid quantitative measurements of relative amounts of H3K9ac and K9me2 in iPSC cells compared to the parental differentiated cells constitute a reliable and convenient criterion to rapidly assess the cell pluripotency potentials and the efficiency of cell reprogramming.

**KEY WORDS:** ESC, iPSC, epigenetic, histone modification

## Introduction

Development of technologies for production of induced pluripotent stem cells (iPSCs) have opened new opportunities to study dynamics of the epigenetic events that underlay pluripotency, differentiation and reprogramming. Re-establishment of the properties of embryonic stem (ES) cell through making iPSC cell generation opens numerous applications in regenerative medicine (Takahashi *et al.*, 2007). Although iPSCs appear as potential alternative to ES cells in various applications, different studies show that they present both genetic and epigenetic abnormalities (Chin *et al.*, 2009, Chin *et al.*, 2010). It appears therefore necessary to define criteria to reliably assess the potentials and quality of iPSC cells. Histone post-translational modifications (PTMs) are at the heart

of many epigenetic mechanisms. Combinations of histone marks create an “epigenetic code” which defines distinct chromatin states and the transcriptional status of the genes (Kouzarides, 2007). Some modifications such as acetylation of histone H3 at lysine 9 (H3K9ac) are correlated with active transcription. This epigenetic mark is abundant on chromatin of ES cells and is known as one of the characteristic features of these cells (Meshorer and Misteli,

*Abbreviations used in this paper:* dif-ESC, differentiated embryonic stem cell; ESC, embryonic stem cell; hESC, human embryonic stem cell; H3K9ac, acetylation of lysine number 9 on histone H3; H3K9me2, di-methylation of lysine 9 on histone H3; H3K4me3, tri-methylation of lysine 4 on histone H3; H3K27me3, tri-methylation of lysine 27 on histone H3; iPSC, induced pluripotent stem cell; PTM, post translational modification.

**\*Address correspondence to:** Maryam Shahhoseini. Royan Institute, No. 2, Hafez St., Banihashem St., Resalat Ave., Tehran-Iran. P. O. Box: 19395-4644. Tel: +98 21 23562737. Fax: +98 21 22310406. E-mail: m.shahhoseini@royaninstitute.org

**Supplementary Material** (four data charts) for this paper is available at: <http://dx.doi.org/10.1387/ijdb.140332ms>

Accepted: 26 May 2016.

2006). Furthermore, the importance of this histone mark during development and mostly in ESC biology has been well substantiated (Hezroni *et al.*, 2011, Wen *et al.*, 2009).

In addition, embryonic stem cells possess bivalent domains that contain coexisting active and repressive histone marks at promoters of developmentally important genes. These bivalent marks refer to chromatin regions bearing two concurrent modifications: tri-methylated lysine 4 and 27 on histone H3 (H3K4me3 and H3K27me3), which are normally associated with gene transcription and gene silencing respectively. Since bivalent marks show reduced levels in differentiated cells compared to ESCs, they appear as important to maintain cell pluripotency (Bernstein *et al.*, 2006, Zhao *et al.*, 2007).

Another critical histone modification is the methylation of histone H3 at lysine 9 (i. e., H3K9me2), which is associated with gene repression in ESCs. This histone mark is found at low levels in ESCs but its occurrence increases during cell differentiation (Wen *et al.*, 2009). Many studies support a critical role for histone H3 lysine 9 (H3K9) methylation in promoting/inhibiting of cell reprogramming and the pluripotency states during iPSC generation (Ang *et al.*, 2011, Singhal *et al.*, 2010).

Interestingly more recent studies point to histone H3 lysine 9 (H3K9) methylation as primary epigenetic determinant acting as a barrier to efficient cell reprogramming both following induced pluripotent stem cell generation and somatic cell nuclear transfer (Chen *et al.*, 2013).

These discoveries allow for the development of specific approaches to assess the pluripotency states of the cells by considering the extent of H3K9 methylation.

In the present study a set of pluripotent and differentiated human cells representing cells going from embryonic state (ESC) into commitment (diff-ESC) and coming back to pluripotency (iPS1), along with human fibroblasts and their reprogrammed counterpart (iPS2), were used to evaluate the ratio of total amounts of H3K9 methylation and acetylation using a quantitative ELISA-based technique. We have also measured following the same method the occurrence of the so-called bivalent H3 modifications: H3K4me and H3K27me. Moreover mRNA expression of several stemness genes (*Nanog*, *Oct4*, *Sox2*) as well as genes indicating fibroblast differentiation (*Vim*, *COL1A1* and *THY1*) were measured quantitatively by qRT-PCR. Using a ChIP-qPCR approach we also tested these four critical histone marks on the regulatory regions of mentioned genes. These investigations allowed us to propose a method to rapidly and reliably assess the cell pluripotency potentials based on the quantitative measurement of H3K9ac/K9me2.


## Results

### ESC differentiation towards fibroblast cells


Fibroblast like cells were derived from ES cells by changing medium and performing passage methods. Morphology of these cells showed the typical spindly shape as normal fibroblast morphology. Immunofluorescence staining depicts the expression of fibroblast marker-Vimentin. The cells were negative for ALP (Alkaline phosphatase), Oct4 and TRA-1-81 (Fig. 1).

### Characterization of iPSC line derived from differentiated ESC

Induced pluripotent stem cell line (iPS1) derived from ESC-



**Fig. 1. Characterization of fibroblast cells differentiated from ESC cells.** (A) Phase contrast image demonstrates typical morphology of fibroblast cells, (B) Immunostaining for Vimentin antigen as an important fibroblast marker, (C) Cells were negative for ALP (Alkaline phosphatase), (D-I) Immunostaining for two stemness markers, Oct-3/4 and TRA-1-81 shows no expression in these cells.




**Fig. 2. Characterization of iPS1 which re-reprogramed from differentiated ESC.** (A) Morphology of iPS1. (B) Higher magnification of iPS1. (C) Expression of ALP (Alkaline phosphatase). (D) Normal Karyotype of cells at passage 12. (E) Expression of Oct-3/4, SSEA-4, TRA-1-60 and TRA-1-81.

fibroblast (dif-ESC) showed normal morphology of hESC colonies as round colonies with definite edge cells with obvious nuclei and thin cytoplasm (Fig. 2A-D). The cells were ALP positive and maintained a normal karyotype with diploid 46 XY karyotype. In addition, immunofluorescent staining showed positive expression of the nuclear marker Oct4 and the surface markers SSEA-4, TRA-1-60 and TRA-1-81 (Fig. 2E).


**Differential expression of stemness and fibroblast genes in iPSC, ESC and differentiated cells**

The relative expression of three stemness genes (*Nanog*, *Oct4* and *Sox2*) and three genes indicating fibroblast differentiation

(*Vim*, *COL1A1* and *THY1*) in all cell lines were quantitatively determined by real-time PCR, using *GAPDH* gene for normalization. As expected, the expression levels of *Nanog*, *Oct4* and *Sox2* were significantly higher in embryonic stem cell (ESC) and in both induced pluripotent stem cells (iPS1 and iPS2) than in dif-ESC and in human fibroblasts (Fig. 3). In contrast, the expression of fibroblast genes (*Vim*, *COL1A1* and *THY1*), in pluripotent cells have showed significantly lower levels than in differentiated cells (Fig. 4) (supplemental data S2). In the case of *COL1A1*, it seemed that in iPS cell lines, especially iPS1, this gene has escaped the global reprogramming and has remained transcribed. In these iPS cells, *Vim* and *THY1* showed a better



**Fig. 3. Relative mRNA expression levels of stemness marker genes in all studied cell lines.** The results are expressed as  $2^{\Delta\Delta CT}$  (mean  $\pm$  SEM). Means labeled with vastly different letters are significantly different in  $p \leq 0.05$  (see supplemental data S2).



**Fig. 4. Relative mRNA expression levels of fibroblast marker genes in all studied cell lines.** The results are expressed as  $2^{-\Delta\Delta CT}$  (mean  $\pm$  SEM). Means labeled with vastly different letters are significantly different in  $p \leq 0.05$  (see supplemental data S2).

reprogramming and present a transcription rate which is closer to that observed in ESCs (Fig. 4).

#### Different levels of histone 3 K9 acetylation and methylation in iPSC, ESC and differentiated cells

Total chromatin extracted from iPSC cell lines, ES cells and differentiated ES cell line were used to evaluate the occurrence H3K9ac and H3K9me2. For this purpose, the chromatin-ELISA method, which allows for a relative quantitative assessment of these histone marks was used. As shown in Fig. 5A, the levels of H3K9ac, in pluripotent cells lines were drastically higher than its level in dif-ESC and fibroblast cells. The same approach was used to measure the relative amounts of H3K9me2 in the same cells. In this case, compared to differentiated cells, we observed low levels of this mark in pluripotent cells (Fig. 5A) (supplemental data S3).

We observed that similarly to H3K9ac, H3K4me3 is higher in pluripotent cells compared to the differentiated cells, while

H3K27me3 showed a reverse situation with higher levels in differentiated cells (Fig. 5B, supplemental data S3).

#### Localized post translational modifications on gene regulatory regions

After evaluation of total levels of H3K9ac/me2 and H3K4/K27 me3 PTMs on chromatin of cells, chromatin immunoprecipitation (ChIP) was used to detect the presence of the mentioned histone modifications on the regulatory regions of specific marker genes indicating stemness states and fibroblast differentiation states respectively in pluripotent and committed cells.

To this aim, after immunoprecipitation of chromatin with H3K9ac, H3K9me2, H3K4me3 and H3K27me3 antibodies, enriched DNA were amplified by using designed primers corresponding to the regulatory regions of the mentioned genes respectively in the considered cell lines. As shown in Fig 6, in all pluripotent cells acetylated H3K9 (H3K9ac) was the dominant modification on regulatory


regions of marker genes of stemness: *Nanog*, *Oct4* and *Sox2*. Likewise, significant high levels of tri-methylated H3K4 were also observed on these regions (see also supplemental data S4).

However, regulatory region of marker genes for fibroblast differentiation (*Vim*, *COL1A1* and *THY1*), were epigenetically marked by hypermethylation of H3K9 and H3K27 in pluripotent cells (Fig. 7).

In fibroblast and differentiated ESC (dif-ESC) the repressive marks of H3K9me2 and H3K27me3 showed significantly higher incorporation than H3K9ac and H3K4me3 at up-stream regions of *Nanog*, *Oct4* and *Sox2* genes (Fig. 6). However, this condition at upstream regions of *Vim*, *COL1A1* and *THY1* was reversed (Fig. 7) (see supplemental data S4).

## Discussion

Embryonic stem cells have specific epigenetic characteristics, which distinguish them from terminally differentiated cells. A re-



**Fig. 5. Total levels of post translational modifications (PTMs) by ELISA method in all studied cell lines.** Relative content of (A) H3K9ac/me2 and (B) bivalent marks of H3K4/27me3 in pluripotent cells of ES, iPS1 and iPS2 as well as dif-ESC and fibroblast cell lines after adjusting for chromatin loading based on H2A reactivity by ELISA. Means labeled with vastly different letters are significantly different in  $p \leq 0.05$  (see supplemental data S3). Asterisk sign shows significant difference of (A) H3K9ac vs. H3K9me2 and (B) H3K4me3 vs. H3K27me3 in  $p \leq 0.05$ .

markable specificity of these cells is their hyper-dynamic chromatin, which has been associated with high levels of histone acetylation (Apostolou and Hochedlinger, 2013, Ikegami *et al.*, 2009, Meshorer *et al.*, 2006). With this respect, histone H3K9 represents a key position. Indeed, H3K9 is a site where competing chemical modifications mediate different functional outputs. Acetylation of H3K9 is associated with both dynamic chromatin and gene transcriptional activity. In contrast, methylation of H3K9 not only blocks further modifications of this specific lysine but also underlays the establishment of heterochromatin and gene transcriptional repression.


Another specificity of ES cells' chromatin is that in these cells, the developmentally important genes are epigenetically distinguished by a unique 'bivalent' mark signature consisting of histone H3

K4 and K27 tri-methylation (Pan *et al.*, 2007). Individually, these histone marks indicate active and repressive genes respectively (Pan *et al.*, 2007).

Since post transcriptional modifications have an undoubted essential role in chromatin dynamicity of cells, it could be a worthwhile way to evaluate the chromatin state of the cells in pluripotency, differentiation and fate determination of ESCs. Here we report an ELISA-based quantitative measurement of total chromatin H3K9ac, H3K9me2 as well as H3K27me3 and H3K4me3 in a specific set of cell lines that has been carefully chosen to reflect different global epigenetic states. Human embryonic stem cells were used as a reference of pluripotent cells and were compared with iPS cells generated from two different sources: differentiated ES cells and fibroblasts.

Our specific simple and quantitative histone PTM measurement approach performed on these cells allowed us to demonstrate a direct relationship between the pluripotency state of the cells and the investigated histone PTMs. More specifically, in pluripotent cells the ratio of total H3K9ac and H3K9me2 was found to indicate the presence of these marks on pluripotency gene promoters as well as the extent of their expression. In contrast to pluripotent cells, in differentiated cells, the ratio of whole chromatin H3K9ac/H3K9me2 is low but interestingly, the active gene promoters are epigenetically distinguished and do not follow the general situation of H3K9 modifications. More precisely, in pluripotent cells, the global amounts of H3K9ac and H3K9me2 mirror the presence of these marks on active gene regulatory regions: the ratio of total H3K9ac/K9me2 is high, so is this ratio on the pluripotency gene promoters (*Nanog*, *OCT4* and *SOX2*). This is not the case of repressed gene promoters (*Vim*, *COL1A1* and *THY1*) in the same cells.

In remarkable contrast, in



**Fig. 6. Chromatin immunoprecipitation (ChIP) analysis of histone modifications on regulatory regions of stemness marker genes.** Incorporated PTMs on regulatory regions of (A,B) *Nonog*, (C,D) *Oct4* and (E,F) *Sox2* in all cell lines (see supplemental data S4).

differentiated cells (fibroblast and dif-ES cells), the ratio of global chromatin K9ac/K9me2 on histone H3 is low, however, promoters of active genes escape this general situation and bear, as expected, higher amounts of H3K9ac compared to H3K9me2.

In agreement with the data presented here, previous studies have demonstrated that H3K9ac is abundant on chromatin of ES cells and is one of the characteristic features of these cells (Meshorer and Misteli, 2006). Additionally, inhibition of histone-deacetylases (HDACs) can enhance gene reactivation in induced pluripotency experiments and promote the reprogramming process (Huangfu et al., 2008).

Histone H3 lysine 9 methylation has been shown in contrast to be an efficient barrier to cell reprogramming both following induced

pluripotent stem cell generation and somatic cell nuclear transfer (Becker et al., 2016).


It is therefore expected that an efficient cell reprogramming of differentiated cells restores high levels of H3K9ac with a corresponding significant reduction in H3K9 methylation. The comparison of the total amount of H3K9ac with that observed in the parental cells and in a reference ES cell line, could therefore be used as a good indicator of the efficiency of cell reprogramming.

In parallel to H3K9ac/H3K9me2, we have also analysed H3K4me3/H3K27me3 at global and loci-specific levels. Our chromatin ELISA tests showed that compared to differentiated cells, higher levels of H3K4me3 could be detected in all three pluripotent cell lines. These results could also be correlated with significant levels

of H3K4me3 upstream of stemness marker genes in pluripotent cells compared to fibroblast and dif-ES cells. A reverse situation was found for H3K27me3, which is more abundant in differentiated cells and marks repressed genes in both ES cells and differentiated cells.

Based on the data presented here, it appears that in pluripotent cells, the promoters of pluripotency genes follow the general situation of whole chromatin H3K9ac/K9me2 as well as total H3K4me3/H3K27me3. This "PTM signature" of chromatin in cells therefore clearly reflects their pluripotency or differentiation state.

Based in these data, we propose that a quantitative measurement of relative amounts of H3K9ac and H3K4me3 compared to H3K9me2 and H3K27me3 in iPS cells with respect to both the parental differentiated cells and a reference ES cell line could constitute a convenient approach to assess the efficiency and success of cell epigenetic reprogramming.



## Materials and Methods

### Cell culture

In this experiment, five human cell lines were studied including an ES cell line (Baharvand et al., 2006, Baharvand et al., 2004) before and

**Fig. 7. Chromatin immunoprecipitation (ChIP) analysis of histone modifications on regulatory regions of fibroblast marker genes.** Incorporated PTMs on regulatory regions of (A,B) THY1, (C,D) Vim and (E,F) COL1A1 in all cell lines (see supplemental data S4).

TABLE 1

## DEFINITION OF THE CELL LINES USED IN THIS STUDY

Cell lines	Definition	References
ESC	Human Embryonic Stem Cell	(Baharvand <i>et al.</i> , 2006, Baharvand <i>et al.</i> , 2004b)
dif-ESC	Differentiated ESC	Unpublished (Hossein Baharvand)
iPS1	Induced pluripotent stem cell reprogrammed from dif-ESC	Unpublished (Hossein Baharvand)
Fibroblast	Human fibroblast	(Mollamohammadi <i>et al.</i> , 2009)
iPS2	Induced pluripotent stem cell reprogrammed from fibroblast	(Mollamohammadi <i>et al.</i> , 2009)

after differentiation to fibroblast (ESC and dif-ESC), and an iPS cell line reprogrammed from these fibroblast cell (iPS1). Accordingly, a human dermal fibroblast cell line as a completely and naturally differentiated cell line was used in this study as well as an iPS line originated from it (iPS2) (Table 1) (Mollamohammadi *et al.*, 2009, Totonchi *et al.*, 2010).

Differentiation process of ESC towards dif-ESC was simply preformed by single cell dissociating of ESCs with TrypLE enzyme (Life Technologies: Cat. No. 12605-028) and seeding single cells on 0.1% gelatin (Sigma-Aldrich: Cat. No. G2500)-coated tissue culture dishes in DMEM medium (Life Technologies: Cat. No. 12800-082) supplemented with 10% FBS (Life Technologies: Cat. No. 16140-071). Passage of the cell line was performed every 3-4 days. Characterization of cells was conducted according to the following parameters: checking of morphology as well as checking of expression of Vimentin (Santa Cruz Biotechnology: Cat. No. sc-32322) as positive marker and Alkaline phosphatase (ALP) (Sigma-Aldrich: Cat. No.86R), Oct-3/4 (Santa Cruz Biotechnology: Cat. No. sc-5279) and TRA-1-81 (Life Technologies: Cat. No. 41-1100) as negative markers. Approved dif-ESC was re-reprogrammed in iPS1 (Totonchi *et al.*, 2010) and standard characterization including morphology, ALP, karyotyping, and expressing Oct-3/4, SSEA-4 (Life Technologies: Cat. No. 41-4000), TRA-1-60 (Life Technologies: Cat. No. 41-1000) and TRA-1-81 were performed. Aforesaid cell lines except fibroblast and dif-ESC were cultured in hESC medium supplemented with 20% knockout serum replacement (Life Technologies: Cat. No. 10828-028), and passage was performed every seven days (Mollamohammadi *et al.*, 2009).

**Karyotype analysis**

For karyotype analysis, at day 5 after culture the cells were treated with 0.66 mM thymidine (Sigma-Aldrich: Cat. No. T1895) for 16 h at 37°C in 5% CO<sub>2</sub>. After being washed, the cells were left for 5 h and then treated with colcemid (Life Technologies: Cat. No. 15212-012) 0.15 mg/ml for 30 min. Isolated human pluripotent stem cells were exposed to 0.075 M KCl at 37°C for 16 min and then were fixed in three consecutive immersions in ice-cold 3:1 methanol to glacial acetic acid and later on dropped onto pre-cleaned chilled slides. Chromosomes were visualized by using standard G-band staining. At least 20 metaphase spreads were screened, and 10 of them were evaluated for chromosomal re-arrangements.

**RNA isolation and quantitative real-time PCR**

RNA isolation and RT-PCR was performed on three biological replicates as described before (Shahhoseini *et al.*, 2010). mRNA quantification was performed in duplicates by quantitative real-time PCR (qRT-PCR) on a StepOnePlus Real-Time PCR System (AB Applied Biosystems) using SYBR Green master mix (AB Applied Biosystems), with designed primers listed in Table 2. qRT-PCR condition was 95°C for 10 minutes, and 40 cycles of 95°C for 15 seconds and 60°C for 60 seconds. Gene expression data were analyzed using  $\Delta\Delta C_t$  quantitative method to estimate relative fold change values.

**Chromatin shearing and soluble chromatin preparation**

To gain nucleosomes as antigens, chromatin was sheared by using a

histone ChIP kit according to manufactured instructions (Diagenode: Cat. No. kch-orgHIS-012). Briefly, each  $1 \times 10^6$  cells were suspended in PBS then 37% formaldehyde (w/v) was added to cells to reach 1% final concentration of formaldehyde and mixed immediately and incubated gently on a shaking platform for 10 minutes at room temperature. To quench the cross-linking reaction of formaldehyde, glycine was added to reach final concentration of 125 mM. Upon being washed with PBS, lysis buffer was added to the pellet of cells and then cells were sonicated for 10 minutes (30"on/30"off, UCD200- Bioruptor sonication system, Diagenode) to get soluble sheared chromatin.

**Chromatin- ELISA analysis**

The chromatin ELISA technique was performed according to Dai and Rasmussen's research (2007) (Dai and Rasmussen, 2007). Briefly, diluted sheared chromatin was coated into 96 well plates (Nunc: 96FMaxisorp, Cat. No. 456537) using coating solution (KPL: Cat. No.50-84-00). After overnight incubation at 4°C, each well was washed four times with 200  $\mu$ L wash solution (KPL Cat. No.50-63-00), then blocked by adding 100  $\mu$ L blocking solution (KPL: Cat. No. 50-84-00) and incubated 1 hour at room temperature (RT). Blocking buffer was aspirated and 50  $\mu$ L of first antibodies (diluted in blocking solution) (see supplemental data S1) were added to each well and incubated at RT for 1-2 hours, then washed as described above. Then, 100  $\mu$ L HRP-conjugated antibodies (supplemental data S1) as secondary antibody (diluted in blocking solution) were added and incubated at RT for 1 hour. After washing, 50  $\mu$ L of 3,3',5,5'-tetra methyl benzidine (TMB) peroxidase substrate (KPL: Cat. No. 50-76-00) was added to each well at RT for 10 minutes. Finally, 50  $\mu$ L, 2 N H<sub>2</sub>SO<sub>4</sub> was added to each well to stop reaction. Optical absorption of plates was read at 450 nm by ELISA-reader (Biotech -ELX 800).

In current study three independent cultures of each cell line were examined as cellular replicates and the soluble chromatin of each replicate was loaded in two or three separate wells for two or three times as technical

TABLE 2

## PRIMER PAIRS USED FOR CHIP REAL-TIME PCR

Gene	Primers (5' → 3')	Annealing temperature(°C)	Product size (bp)
<b>qRT-PCR primers</b>			
NANOG	F: AAAGTCTTAAAGCTGCCTTAAC R: CAGTCGGATGCTTCAAAG	60	130
OCT4	F: GTTCTTCATTCTACTAAGGAAGG R: CAAGAGCATCATTGAACCTCAC	60	101
SOX2	F:GGGAAATGGGAGGGGTGCAAAAGAGG R:TTGCGTGAGTGTGGATGGGATTGGTG	60	151
VIM	F:GGCTCGTCACCTTCGTGAAT R:GAGAAATCCTGCTCTCTCTCGC	60	110
THY1	F:TAGTGAAGGCGGATAAGTAGAGG R: ACCCGTGAGACAAAGAAGC	60	124
COL1A1	F:CCTGTCTGCTTCTCTGTAAC R:ATGTTCCGGTTGGTCAAAGATAAA	60	211
<b>ChIP Real-time PCR Primer</b>			
NANOG	F: AATTCACAAGGGTGGGTGAG R:TAACATGAGGCAACCAGCTC	60	133
OCT4	F: GTTGGGGAGCAGGAAGCA R: GGGGAGCTCTAACCCCTAAA	60	83
SOX2	F: TCGCTAGAAACCCATTATTTC R: CTGCCCTTGACAACCTCTG	60	93
VIM	F: CGAGTTTCTCTTTCCACC R: ACTTCTGCAGCCTTTGGA	60	187
THY1	F: GGCTTTAACCTTTCTTCTGAC R: AAGGAAATGTGGAGCGT	60	106
COL1A1	F: GGGCCCTTTTATACTGTCC R: TCTCCATTCCAACCTCCAAA	60	132

NANOG (Nanog homeobox), OCT4 (POU class 5 homeobox 1), SOX2 (SRF box2), VIM (Vimentin), THY1 (Thy-1 cell surface antigen), COL1A1 (Collagen, type I, alpha 1)

replicates. After subtracting of background from each well, values were normalized by dividing them into loading-controlled values. At least, values were expressed as means $\pm$ SEM of 3 separate experiments.

#### Chromatin immunoprecipitation (ChIP)-real time PCR analysis

Chromatin immunoprecipitation (ChIP) experiments were conducted as previously described on three biological replicates (Favaedi et al., 2012). For each immunoprecipitation reaction  $1 \times 10^5$  cells was used. Recovered DNA from precipitation fractions and total chromatin input were applied for quantitative real-time PCR with designed primers for regulatory regions of 6 genes involved in stemness and fibroblast differentiation (Table 2). Real-time PCR was performed on a step one plus Real-Time PCR System (Applied Biosystems) by using SYBR Green PCR master mix (Applied Biosystems: Cat. No. 4367659). PCR condition was 95°C for 10 min; and 40 cycles of 95°C for 15 s, 60°C for 45 s. The results are expressed as %input which means percentage of enriched DNA associated with given immunoprecipitated histone modifications relative to a 1/100 dilution of input chromatin (mean  $\pm$  SEM; triplicate experiment).

#### Statistical analysis

Values are expressed as means $\pm$ SEM of three separate experiments. All data was analyzed by using One-way ANOVA and paired samples T-test, and differences were considered statistically significant at  $p \leq 0.05$  (see supplemental data S2 and S3).

#### Acknowledgments

We wish to thank Dr. Saadi.Khochbin for his worthwhile guidance and Farideh Moeinvaziri for her kind help. The authors would like to dedicate this paper to the memory of Dr. Saeid Kazemi Ashtiani, the late founder of Royan Institute. This project was financially supported by the Royan Institute (Grant No. 173).

#### References

- ANG, Y.-S., TSAI, S.-Y., LEE, D.-F., MONK, J., SU, J., RATNAKUMAR, K., DING, J., GE, Y., DARR, H. AND CHANG, B. (2011). WDR5 MEDIATES SELF-RENEWAL AND REPROGRAMMING VIA THE EMBRYONIC STEM CELL CORE TRANSCRIPTIONAL NETWORK. *Cell* 145: 183-197.
- APOSTOLOU, E. AND HOCHEDLINGER, K. (2013). CHROMATIN DYNAMICS DURING CELLULAR REPROGRAMMING. *Nature* 502: 462-471.
- BAHARVAND, H., ASHTIANI, S.K., TAEI, A., MASSUMI, M., VALOJERDI, M.R., YAZDI, P.E., MORADI, S.Z. AND FARROKHI, A. (2006). GENERATION OF NEW HUMAN EMBRYONIC STEM CELL LINES WITH DIPLOID AND TRIPLOID KARYOTYPES. *Dev Growth Differ* 48: 117-128.
- BAHARVAND, H., ASHTIANI, S.K., VALOJERDI, M.R., SHAHVERDI, A., TAEI, A. AND SABOUR, D. (2004). ESTABLISHMENT AND *IN VITRO* DIFFERENTIATION OF A NEW EMBRYONIC STEM CELL LINE FROM HUMAN BLASTOCYST. *Differ* 72: 224-229.
- BECKER, J.S., NICETTO, D., ZARET, K.S. (2016). H3K9me3-Dependent Heterochromatin: Barrier to Cell Fate Changes. *Trends Genet.* 32: 29-41.
- BERNSTEIN, B.E., MIKKELSEN, T.S., XIE, X., KAMAL, M., HUEBERT, D.J., CUFF, J., FRY, B., MEISSNER, A., WERNIG, M. and PLATH, K. (2006). A bivalent chromatin structure marks key developmental genes in embryonic stem cells. *Cell* 125: 315-326.
- CHEN, J., LIU, H., LIU, J., QI, J., WEI, B., YANG, J., LIANG, H., CHEN, Y., CHEN, J. and WU, Y. (2013). H3K9 methylation is a barrier during somatic cell reprogramming into iPSCs. *Nat Genet* 45: 34-42.
- CHIN, M.H., MASON, M.J., XIE, W., VOLINIA, S., SINGER, M., PETERSON, C., AMBARTSUMYAN, G., AIMIUWU, O., RICHTER, L. and ZHANG, J. (2009). Induced pluripotent stem cells and embryonic stem cells are distinguished by gene expression signatures. *Cell Stem Cell* 5: 111-123.
- CHIN, M.H., PELLEGRINI, M., PLATH, K. and LOWRY, W.E. (2010). Molecular analyses of human induced pluripotent stem cells and embryonic stem cells. *Cell Stem Cell* 7: 263-269.
- DAI, B. and RASMUSSEN, T.P. (2007). Global epiproteomic signatures distinguish embryonic stem cells from differentiated cells. *Stem Cells* 25: 2567-2574.
- FAVAEDI, R., SHAHHOSEINI, M., AKHOOND, M.R. (2012). Comparative epigenetic analysis of Oct4 regulatory region in RA-induced differentiated NT2 cells under adherent and non-adherent culture conditions. *Mol Cell Biochem* 363: 129-134.
- HEZRONI, H., TZCHORI, I., DAVIDI, A., MATTOU, A., BIRAN, A., NISSIM-RAFINIA, M., WESTPHAL, H. and MESHORER, E. (2011). H3K9 histone acetylation predicts pluripotency and reprogramming capacity of ES cells. *Nucleus* 2: 300-309.
- HUANGFU, D., MAEHR, R., GUO, W., EIJKELENBOOM, A., SNITOW, M., CHEN, A.E. and MELTON, D.A. (2008). Induction of pluripotent stem cells by defined factors is greatly improved by small-molecule compounds. *Nat Biotechnol* 26: 795-797.
- IKEGAMI, K., OHGANE, J., TANAKA, S., YAGI, S. and SHIOTA, K. (2009). Interplay between DNA methylation, histone modification and chromatin remodeling in stem cells and during development. *Int J Dev Biol* 53: 203-214.
- KOUZARIDES, T. (2007). Chromatin modifications and their function. *Cell* 128: 693-705.
- MESHORER, E. and MISTELI, T. (2006). Chromatin in pluripotent embryonic stem cells and differentiation. *Nat Rev Mol Cell Biol* 7: 540-546.
- MESHORER, E., YELLAJOSHULA, D., GEORGE, E., SCAMBLER, P.J., BROWN, D.T. and MISTELI, T. (2006). Hyperdynamic plasticity of chromatin proteins in pluripotent embryonic stem cells. *Dev Cell* 10: 105-116.
- MOLLAHAMMADI, S., TAEI, A., PAKZAD, M., TOTONCHI, M., SEIFINEJAD, A., MASOUDI, N. and BAHARVAND, H. (2009). A simple and efficient cryopreservation method for feeder-free dissociated human induced pluripotent stem cells and human embryonic stem cells. *Hum Reprod* 24: 2468-2476.
- PAN, G., TIAN, S., NIE, J., YANG, C., RUOTTI, V., WEI, H., JONSDOTTIR, G.A., STEWART, R., THOMSON, J.A. (2007). Whole-genome analysis of histone H3lysine 4 and lysine 27 methylation in human embryonic stem cells. *Cell Stem Cell* 1: 299-312.
- SINGHAL, N., GRAUMANN, J., WU, G., ARAÚZO-BRAGO, M.J., HAN, D.W., GREBER, B., GENTILE, L., MANN, M. and SCHÖLER, H.R. (2010). Chromatin-remodeling components of the BAF complex facilitate reprogramming. *Cell* 141: 943-955.
- TAKAHASHI, K., TANABE, K., OHNUKI, M., NARITA, M., ICHISAKA, T., TOMODA, K. and YAMANAKA, S. (2007). Induction of pluripotent stem cells from adult human fibroblasts by defined factors. *Cell* 131: 861-872.
- TOTONCHI, M., TAEI, A., SEIFINEJAD, A., TABEBORDBAR, M., RASSOULI, H., FARROKHI, A., GOURABI, H., AGHDAMI, N., HOSSEINI-SALEKDEH, G. and BAHARVAND, H. (2010). Feeder- and serum-free establishment and expansion of human induced pluripotent stem cells. *Int J Dev Biol* 54: 877.
- WEN, B., WU, H., SHINKAI, Y., IRIZARRY, R.A. and FEINBERG, A.P. (2009). Large histone H3 lysine 9 dimethylated chromatin blocks distinguish differentiated from embryonic stem cells. *Nat Genet* 41: 246-250.
- ZHAO, X.D., HAN, X., CHEW, J.L., LIU, J., CHIU, K.P., CHOO, A., ORLOV, Y.L., SUNG, W.-K., SHAHAB, A. and KUZNETSOV, V.A. (2007). Whole-genome mapping of histone H3 Lys4 and 27 trimethylations reveals distinct genomic compartments in human embryonic stem cells. *Cell Stem Cell* 1: 286-298.



**Further Related Reading, published previously in the *Int. J. Dev. Biol.***

**Bone morphogenetic protein 4 promotes craniofacial neural crest induction from human pluripotent stem cells**

Sumiyo Mimura, Mika Suga, Kaori Okada, Masaki Kinehara, Hiroki Nikawa and Miho K. Furue  
Int. J. Dev. Biol. (2016) 60: 21-28  
<http://dx.doi.org/10.1387/ijdb.160040mk>

**Two decades of reproductive biomedicine and stem cell biology in Iran: the Royan Institute**

Sophie Rousseaux  
Int. J. Dev. Biol. (2014) 58: 643-647  
<http://dx.doi.org/10.1387/ijdb.140245sr>

**Generation of pluripotent stem cells via protein transduction**

Xia Li, Pengfei Zhang, Chao Wei and Yunhai Zhang  
Int. J. Dev. Biol. (2014) 58: 21-27  
<http://dx.doi.org/10.1387/ijdb.140007XL>

**Epigenetic features of testicular germ cell tumours in relation to epigenetic characteristics of foetal germ cells**

Dina G. Kristensen, Niels E. Skakkebaek, Ewa Rajpert-De Meyts and Kristian Almstrup  
Int. J. Dev. Biol. (2013) 57: 309-317  
<http://dx.doi.org/10.1387/ijdb.130142ka>

**Signaling pathways during maintenance and definitive endoderm differentiation of embryonic stem cells**

Lina Sui, Luc Bouwens and Josu e K. Mfopou  
Int. J. Dev. Biol. (2013) 57: 1-12  
<http://dx.doi.org/10.1387/ijdb.120115ls>

**5 yr ISI Impact Factor (2013) = 2.879**

