

doi: 10.1387/ijdb.170037af


SUPPLEMENTARY MATERIAL

corresponding to:

**Expression of the ALK1 family of type I BMP/ADMP receptors
during gastrula stages in *Xenopus* embryos**

AVI LEIBOVICH, HERBERT STEINBEISER and ABRAHAM FAINSOD*


*Address correspondence to: Abraham Fainsod, Department of Developmental Biology and Cancer Research, Institute for Medical Research Israel-Canada, Faculty of Medicine, Hebrew University, Jerusalem 9112102, Israel. Tel: +972-2-6758157. Fax: +972-2-6757482. E-mail: abrahamf@ekmd.huji.ac.il

 <http://orcid.org/0000-0002-6368-0416>

Full text for this paper is available at: <http://dx.doi.org/10.1387/ijdb.170037af>


A

ALK1 (ACVRL1)



B

ALK2 (ACVR1)



Supplementary Fig. S1. The ALK1 (ACVRL1) and ALK2 (ACVR1) receptors in *Xenopus laevis*. Protein sequence comparison of the two (.S and.L) ALK1 (A) and ALK2 (B) receptors in the *Xenopus laevis* genome. Comparisons were performed to their counterparts from *Xenopus tropicalis*, chicken and human. Amino acids in dark gray represent identical residues while light gray shading identifies similar amino acids. Gaps are marked with dashes.